# **PCT**

#### WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: A61K 31/07, 31/19, 31/35, 31/44, 31/075 A1

(11) International Publication Number:

WO 00/06144

(43) International Publication Date:

10 February 2000 (10.02.00)

(21) International Application Number:

PCT/US99/16866

(22) International Filing Date:

26 July 1999 (26.07.99)

(30) Priority Data:

09/123,484

27 July 1998 (27.07.98)

US

(71)(72) Applicant and Inventor: MURAD, Howard [US/US]; Penthouse 11, 4265 Marina City Drive, Marina del Rey, CA 90292 (US).

(74) Agents: FANUCCI, Allan, A. et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

### Published

With international search report.

(54) Title: PHARMACEUTICAL COMPOSITIONS AND METHODS FOR MANAGING SCALP CONDITIONS

### (57) Abstract

This application relates to a pharmaceutical composition for the prevention, treatment, and management of scalp conditions, such as dandruff, seborrheic dermatitis, psoriasis, folliculitis, and hair thinning including a therapeutically effective amount of an acidic component of a hydroxy acid or tannic acid, or a pharmaceutically acceptable salt thereof. A preferred anti-dandruff composition and method of managing dandruff includes a therapeutically effective amount of the acid component, a vitamin A component, and an anti-growth agent. A preferred anti-hair thinning composition and method of managing thinning hair includes a therapeutically effective amount of the acidic component, a niacin component present in an amount sufficient to locally increase blood circulation, and a  $5-\alpha$  reductase inhibitor. The invention also relates to a method of treating chemically processed hair by administering to a patient an amount of an acidic component of a hydroxy acid or tannic acid, or a pharmaceutically acceptable salt thereof, in an amount sufficient to essentially close the cuticle and inhibit modification of the chemically processed hair.

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	МD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	MŁ	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Tceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# PHARMACEUTICAL COMPOSITIONS AND METHODS FOR MANAGING SCALP CONDITIONS

## TECHNICAL FIELD

20

35

This application relates to pharmaceutical compositions, as well as methods, to normalize skin for the prevention, treatment, and management of scalp conditions.

# 10 BACKGROUND OF THE INVENTION

Human skin is a composite material of the epidermis and the dermis. The topmost part of the epidermis is the stratum corneum. This layer is the stiffest layer of the skin, as well as the one most affected by the surrounding environment. Below the stratum corneum is the internal portion of the epidermis. Below the epidermis, the topmost layer of the dermis is the papillary dermis, which is made of relatively loose connective tissues that define the micro-relief of the skin. The reticular dermis, disposed beneath the papillary dermis, is tight, connective tissue that is spatially organized. The reticular dermis is also associated with coarse wrinkles. At the bottom of the dermis lies the subcutaneous layer.

The principal functions of the skin include protection, excretion, secretion, absorption, thermoregulation, pigmentogenesis, accumulation, sensory perception, and regulation of immunological processes. These functions are detrimentally affected by, for example, dryness, yeast, and structural changes in the skin, such as due to aging and excessive sun exposure.

Many hair care products tend to include alkaline components, which tend to open pores in the scalp. While these products may improve or alter hair appearance, they may exacerbate various scalp conditions, such as dandruff, psoriasis, and seborrhea.

Various pharmaceuticals that frequently contain one or more alpha hydroxy acids, such as glycolic acid, have been used to prevent and treat certain cellular, skin, hair and other conditions. For example, U.S. Patent No. 4,668,509 discloses polythioalkanecarboxylic anionic products and their preparation and use in cosmetic compositions and hair treatment compositions, such as shampoos.

U.S. Patent No. 4,814,166 discloses polyanionic oligomer compounds suitable for use in keratin fiber treatment, such as hair, which may be administered in a shampoo. These compounds are alleged to hold hair with suppleness and without significant hardening of the hair.

U.S. Patent No. 5,344,971 discloses mercapto acids, such as thiolactic acid (2-mercaptopropionic acid), that are used as reducing agents for the permanent reshaping of hair or for depilatory milks and creams.

An article entitled "Hydroxy Acids and Skin Aging" discloses the use of hydroxy and other acids as skin peels and emollients that can moisturize, stimulate, and exfoliate the skin. [Smith, W., Soap/Cosmetics/Chemical Specialties, pp. 54-58, 76, Sept., 1993.]. A study was conducted with hydroxy, keto, carboxylic, or dicarboxylic acids, including glycolic and salicylic acids, to determine long-term rejuvenating benefits. The study noted that higher pH formulations resulted in less stimulatory activity and lower irritation, and concluded that various non-hydroxy acids would be expected to deliver long-term rejuvenating benefits. *Id.* at 58.

One publication, WO 95/33438, discloses skin or hair care products having an agent acting cosmetically on the hair or skin, such as thioglycolic acid preferably with an ammonium salt. The agent is absorbed in a fibrous material containing amino groups, such as cellulose based-fiber containing polysilicic acid.

U.S. Patent No. 5,422,370 discloses methods of using hydroxyacid or related compounds, such as alpha 2-hydroxypropanoic acid (lactic acid), for the treatment of wrinkles. These compounds are also disclosed to be effective for enhancing the topical effects of other cosmetic and pharmaceutical agents for treatment of conditions such as dry skin, ichthyosis, eczema, palmar and plantar hyperkeratoses, dandruff, acne, pruritis, psoriasis, Darier's disease, lichen simplex chronicus, and warts. Similarly, U.S. Patent No. 5,547,988 discloses methods for reducing the

warts. Similarly, U.S. Patent No. 5,547,988 discloses methods for reducing the appearance of skin changes associated with aging, such as wrinkles, by topically applying a compound of glycolic acid, lactic acid, citric acid, or a salt thereof.

U.S. Patent No. 5,587,149 discloses improved stable emulsions of polyethylene glycol-in-oil for topical application to the skin that contain one or more water soluble active ingredients, such as Vitamin C, glycolic acid, and the like.

PCT/US99/16866 WO 00/06144

Despite these references, there is still a need for pharmaceutical compositions and methods for the prevention, treatment, and management of scalp conditions, such as dandruff, psoriasis, and seborrhea. The present invention advantageously provides pharmaceutical compositions, as well as methods for prevention and treatment, by administering such compositions to repair and normalize the scalp for the prevention and treatment of scalp conditions.

## SUMMARY OF THE INVENTION

10

The present invention relates to an anti-dandruff pharmaceutical composition for administration to a scalp including an acidic component having a hydroxy acid or tannic acid, or a pharmaceutically acceptable salt thereof, in an amount sufficient to exfoliate at least a portion of the scalp, a vitamin A component in an amount sufficient to inhibit 15 overproduction of stratum corneum on the scalp, and an anti-growth agent to inhibit fungus, yeast, bacteria, or a mixture thereof adjacent the scalp. The compositions preferably include a pharmaceutically acceptable carrier or excipient.

In one embodiment, the acidic component includes an alpha-hydroxy acid, beta-hydroxy acid or tannic acid, the vitamin A component includes retinyl palmitate, and the anti-growth agent includes clotrimazole. In a preferred embodiment, the acidic component includes at least one of glycolic acid, lactic acid, citric acid, salicylic acid, or tannic acid. In another embodiment, the acidic component is present in an amount from about 0.1 to 8 weight percent, the vitamin A component is present in an amount from about 0.01 to 1 weight percent, and the anti-growth agent is present in an amount from about 0.1 to 1.5 weight percent, of the composition. In a preferred embodiment, the pharmaceutical composition further includes at least one of a surfactant, a stabilizer. a preservative, a moisturizer, anti-inflammatory agent, anti-oxidant, and a coloring 30 agent, which together may be present in an amount from about 10.1 to 99.1 weight percent of the composition. Also encompassed by the invention are a shampoo, cream. or gel including the pharmaceutical composition described above.

The invention also relates to an anti-hair thinning pharmaceutical 35 composition for administration to a scalp having an acidic component including a hydroxy acid or tannic acid in an amount sufficient to exfoliate at least a portion of the

PCT/US99/16866 WO 00/06144

scalp, or a pharmaceutically acceptable salt thereof, a niacin component present in an amount sufficient to locally increase blood circulation; and a  $5-\alpha$  reductase inhibitor in an amount sufficient to inhibit conversion of testosterone to dihydro-testosterone. The compositions preferably include a pharmaceutically acceptable carrier or excipient.

In one embodiment, the acidic component includes an alpha-hydroxy acid, beta-hydroxy acid. or tannic acid, the niacin component includes nicotinate, and the  $5-\alpha$ reductase inhibitor includes at least one of finasteride or Saw Palmetto Extract. In a preferred embodiment, the acidic component includes glycolic acid, lactic acid, citric acid, salicylic acid, or tannic acid. In another embodiment, the acidic component is present in an amount from about 0.1 to 8 weight percent, the niacin component is present in an amount from about 0.01 to 1 weight percent, and the 5- $\alpha$  reductase inhibitor is present in an amount from about 0.1 to 1 weight percent, of the 15 composition. In a preferred embodiment, the composition includes at least one of a surfactant, a stabilizer, a preservative, a moisturizer, anti-inflammatory agent, antioxidant, and a coloring agent, which together may be present in an amount from about 10.1 to 99.1 weight percent of the composition. In a preferred embodiment, the acidic component and the niacin component are adapted for topical administration and the 5- $\alpha$ reductase inhibitor is adapted for oral administration. The invention also encompasses a shampoo, cream, or gel including the pharmaceutical composition described above.

The invention also relates to a method of managing a scalp condition which includes administering to a patient a therapeutically effective amount of an acidic component having a hydroxy acid or tannic acid, or a pharmaceutically acceptable salt thereof, a vitamin A component, and an anti-growth agent to inhibit fungus, yeast, bacteria, or a mixture thereof that may be present adjacent the scalp. In a preferred embodiment, the scalp condition treated is at least one of dandruff, seborrheic 30 dermatitis, psoriasis, folliculitis, or other infectious or scaling conditions.

25

In a preferred embodiment, the active ingredients are administered topically. In another embodiment, the administering is by at least one of a shampoo, aerosol spray, gel, paste, cream, lotion, sponge, emulsion, or ointment. In a preferred 35 embodiment, from about 1 mg to 10,000 mg of the acidic component, vitamin A component, and anti-growth agent together are administered. In another embodiment,

the acidic component, vitamin A component, and anti-growth agent are administered concurrently. In another embodiment, the acidic component, vitamin A component, and anti-growth agent are administered concurrently with at least one additional pharmaceutical composition for the prevention or treatment of a scalp condition. In another embodiment, the method further includes administering at least one of a surfactant, stabilizer, preservative, moisturizer, anti-inflammatory agent, anti-oxidant, or coloring agent. In a preferred embodiment, the acidic component includes an alphahydroxy acid or tannic acid, the vitamin A component includes retinyl palmitate, and the anti-growth agent includes clotrimazole.

The invention also relates to a method of managing hair thinning which includes administering to a patient a therapeutically effective amount of an acidic component having a hydroxy acid or tannic acid, or a pharmaceutically acceptable salt thereof, a niacin component present in an amount sufficient to locally increase blood supply, and a 5-α reductase inhibitor.

In a preferred embodiment, the active ingredients are administered topically. In a more preferred embodiment, the acidic component and niacin component are topically administered and the  $5-\alpha$  reductase inhibitor is orally administered. In another embodiment, the administering is by at least one of a shampoo, aerosol spray, gel, paste, cream, lotion, sponge, emulsion, or ointment. In a preferred embodiment, from about 1 mg to 10,000 mg of the acidic component, niacin component, and  $5-\alpha$ reductase inhibitor are administered. In another embodiment, the acidic component, niacin component, and  $5-\alpha$  reductase inhibitor are administered concurrently. In yet another embodiment, the acidic component, niacin component, and  $5-\alpha$  reductase inhibitor are administered concurrently with at least one additional pharmaceutical composition for the prevention or treatment of a scalp condition. In another 30 embodiment, the method further includes administering at least one of a surfactant, stabilizer, preservative, moisturizer, anti-inflammatory agent, anti-oxidant, or coloring agent. In a preferred embodiment, the acidic component includes an alpha-hydroxy acid or tannic acid, the niacin component includes nicotinate, and the 5- $\alpha$  reductase inhibitor 35 includes at least one of finasteride or Saw Palmetto Extract.

The invention further relates to a method of treating chemically processed hair having a plurality of cuticle by administering to a patient an amount of an acidic component including a hydroxy acid or tannic acid, or a pharmaceutically acceptable salt thereof, in an amount sufficient to essentially close the cuticle.

In a preferred embodiment, the administering is topically applied. In another embodiment, the administering is by at least one of a shampoo, aerosol spray, gel, paste, cream, lotion, sponge, emulsion, or ointment. In a preferred embodiment, from about 1 mg to 10,000 mg of the acidic component is administered. In another embodiment, the acidic component is administered concurrently with at least one additional pharmaceutical composition for the prevention or treatment of a scalp condition. In a preferred embodiment, the method further includes administering at least one of a surfactant, stabilizer, preservative, moisturizer, anti-inflammatory agent, anti-oxidant, or coloring agent. In another preferred embodiment, the acidic component includes an alpha-hydroxy acid or tannic acid.

The ranges of the components of the pharmaceutical composition may vary, but the active ingredients should be understood to add to 100 weight percent of the active pharmaceutical composition.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

20

A formulation for the prevention, treatment, and management of scalp conditions, such as dandruff, psoriasis, and seborrhea, has now been discovered. Moreover, the management of the scalp may advantageously be accomplished by the administration of the pharmaceutical composition of the present invention.

Methods for administering the compositions herein are also encompassed by the invention. Such methods include the prevention, treatment, or management of:

30 dandruff, eczema, hair thinning, irritation, psoriasis, folliculitis, scaling, seborrhea or seborrheic dermatitis, all while reducing environmental damage to the hair and scalp and maintaining color in chemically processed hair as advantageous benefits. The compositions may be prepared in high concentrations for administration by professionals in a salon, as well as lower concentrations that are safer for home use. The methods of the present invention also include administering mono- or poly-hydroxy or tannic acids

for treating chemically processed hair, which advantageously closes or essentially closes the hair cuticle. The closing of the cuticle helps "lock in" hair color to prevent fading or washout of the chemically processed hair.

5

The pharmaceutical composition includes the combination of a number of different components that interact to provide the desired improvements to the skin. The compositions include an acidic component including one or more mono- or poly-hydroxy acids or tannic acid, a mixture thereof, or a pharmaceutically acceptable salt or ester thereof. One of ordinary skill in the art will be readily able to select and prepare suitable mono- or poly-hydroxy acids for use in the composition of the invention, for example, alkyl hydroxycarboxylic acids, aralkyl and aryl hydroxycarboxylic acids, polyhydroxy-carboxylic acids, and hydroxy-polycarboxylic acids. To assist one of ordinary skill in the art, some preferred mono- or poly-hydroxy acids include: 2-15 hydroxyacetic acid (glycolic acid); 2-hydroxypropanoic acid (lactic acid); 2-methyl 2hydroxypropanoic acid; 2-hydroxybutanoic acid; phenyl 2-hydroxyacetic acid; phenyl 2methyl 2-hydroxyacetic acid; 3-phenyl 2-hydroxyacetic acid; 2,3-dihydroxypropanoic acid; 2,3,4-trihydroxybutanoic acid; 2,3,4,5,6-pentahydroxyhexanoic acid;

2-hydroxydodecanoic acid; 2,3,4,5-tetrahydroxypentanoic acid; 2,3,4,5,6,7-hexahydroxyheptanoic acid; diphenyl 2-hydroxyacetic acid; 4-hydroxymandelic acid; 4-chloromandelic acid; 3-hydroxybutanoic acid; 4-hydroxybutanoic acid; 2-hydroxyhexanoic acid; 5-hydroxydodecanoic acid; 12-hydroxydodecanoic acid; 10-hydroxydecanoic acid; 16-hydroxyhexadecanoic acid;

2-hydroxy-3-methylbutanoic acid; 2-hydroxy-4-methylpentanoic acid; 3-hydroxy-4methoxymandelic acid; 4-hydroxy-3-methoxymandelic acid; 2-hydroxy-2-methylbutanoic acid; 3-(2-hydroxphenyl) lactic acid; 3-(4-hydroxyphenyl) lactic acid; hexahydromandelic acid; 3-hydroxy-3-methylpentanoic acid; 4-hydroxydecanoic acid;

30 5-hydroxydecanoic acid; aleuritic acid; 2-hydroxypropanedioic acid; 2-hydroxybutanedioic acid; erythraric acid; threaric acid; arabiraric acid; ribaric acid; xylaric acid; lyxaric acid; glucaric acid; galactaric acid; mannaric acid; gularic acid; allaric acid; altraric acid; idaric acid; talaric acid; 2-hydroxy-2-methylbutanediodic acid; 35 citric acid, isocitric acid, agaricic acid, quinic acid, glucoronic acid, glucoronolactone, galactoronic acid, galactoronolactone, uronic acids, uronolactones, ascorbic acid,

dihydroascorbic acid, dihydroxytartaric acid, tropic acid, ribonolactone, gluconolactone. galactonolactone, gulonolactone, mannonolactone, citramalic acid; pyruvic acid, hydyroxypyruvic acid, hydroxypyruvic acid phosphate and esters thereof; methyl pyruvate, ethyl pyruvate, propyl pyruvate, isopropyl pyruvate; phenyl pyruvic acid and esters thereof; methyl phenyl pyruvate, ethyl phenyl pyruvate, propyl phenyl pyruvate; formyl formic acid and esters thereof; methyl formyl formate, ethyl formyl formate. propyl formyl formate; benzoyl formic acid and esters thereof; methyl benzoyl formate. ethyl benzoyl formate and propyl benzoyl formate; 4-hydroxybenzoyl formic acid and esters thereof; 4-hydroxyphenyl pyruvic acid and esters thereof; and 2-hydroxyphenyl pyruvic acid and esters thereof. The hydroxy acids are more preferably selected from one or more alpha-hydroxy acids or beta-hydroxy acids, most preferably from glycolic. lactic, citric, or salicylic acid. It should be understood that one or more derivatives of 15 the above acidic component, such as esters or lactones thereof, are also suitably used. One of ordinary skill in the art will also understand that various hydroxy acids described in U.S. Patent Nos. 5,547,988 and 5,422,370 are also suitable for use in the compositions and methods of the invention. The acidic component is present in the composition and methods in an amount sufficient to exfoliate, i.e., remove dead or dying skin cells, from at least a portion of the scalp. The acidic component is typically present in an amount from about 0.1 to 8 weight percent, preferably about 0.5 to 5 weight percent, more preferably from about 1 to 3 weight percent of the composition. For example, the acidic component may be from about 0.2 to 10 weight percent GLYPURE, which contains 70 weight percent glycolic acid, in combination with about 0.1 to 0.3 weight percent salicylic acid.

Compositions and methods for managing dandruff, seborrheic dermatitis, psoriasis, folliculitis, and other scaling or infectious scalp conditions, also include a vitamin A component and an anti-growth agent. The vitamin A component preferably is retinyl palmitate. The vitamin A component is present in an amount sufficient to normalize epidermal growth by inhibiting overproduction of the stratum corneum. The vitamin A component is typically present in an amount from about 0.01 to 1 weight percent, preferably 0.05 to 0.8 weight percent, and more preferably 0.1 to 0.5 weight

percent of the composition. Vitamin A is toxic at high levels, such that no more than 400,000 IU should be cumulatively ingested per day for greater than six months.

Any pharmaceutically acceptable anti-growth agent may be used,

preferably an antifungal agent, antiyeast agent, or antibacterial agent, more preferably
triclosan, clotrimazole, and the like, and mixtures thereof. The anti-growth agent is
typically present in an amount from about 0.1 to 1.5 weight percent, preferably from
about 0.3 to 1.2 weight percent, and more preferably from about 0.5 to 1 weight
percent of the composition. The anti-growth agent may inhibit the fungus *Pityrosporum*ovale, which tends to be present in patients having dandruff or seborrheic dermatitis.
Together, the acidic component, Vitamin A component, and anti-growth agent facilitate
exfoliation of dead skin, facilitate hydration of the scalp, and inhibit the presence of
yeast or fungus, such as all of which assist in the management of dandruff, psoriasis,
folliculitis, seborrheic dermatitis, and other inflammatory or scaling conditions of the
scalp.

Compositions and methods for the management of thinning hair include a niacin component and a 5-α reductase inhibitor, in addition to the acidic component.

The niacin component (vitamin B<sub>3</sub>) may be any pharmaceutically acceptable niacin component, preferably niacinamide or nicotinate, more preferably nicotinate. The niacin component should be present in an amount sufficient to facilitate improved blood circulation in the scalp, which may help inhibit hair loss. Thus, the niacin component is typically present in an amount from about 0.01 to 1 weight percent, preferably from about 0.05 to 0.8 weight percent, and more preferably from about 0.1 to 0.5 weight percent of the composition.

Any pharmaceutically acceptable 5- $\alpha$  reductase inhibitor may be used in the anti-hair thinning compositions, such as finasteride, Saw Palmetto Extract (Serenoa repens), MK-386, LY191704, turosteride, PNU 157706, FK-143, GG-745, ONO-9302, and the like, and mixtures thereof. A preferred 5- $\alpha$  reductase inhibitor is Saw Palmetto Extract (Serenoa Serrulata). The 5- $\alpha$  reductase inhibitor is typically present in an amount sufficient to inhibit conversion of testosterone in the scalp to dihydrotestosterone, the latter of which is believed to increase hair thinning. The 5- $\alpha$  reductase inhibitor is typically present in an amount from about 0.01 to 1 weight percent,

preferably about 0.05 to 0.5 weight percent, and more preferably about 0.08 to 0.2 weight percent of the composition. The hydroxy acid, niacin component, and  $5-\alpha$  reductase inhibitor together facilitate exfoliation of dead skin, improve blood circulation, and ultimately inhibit and/or reduce hair loss.

In a preferred embodiment, the compositions all contain one or more surfactants, stabilizers, preservatives, moisturizers, coloring agents, anti-inflammatory agents, anti-oxidants, water, and the like, and mixtures thereof. The water used is preferably deionized water. It should be understood that water includes the remainder of a given composition after other ingredients are determined. Although any pharmaceutically acceptable surfactant, stabilizer, preservative, moisturizer, or coloring agent may be used, certain compounds or mixtures are preferred as discussed below.

Preferred surfactants, including both the foaming and non-foaming type, include sodium laureth sulfate, disodium cocoamphodiacetate, and the like, and mixtures thereof. The surfactant component may be present in an amount from about 10 to 90 weight percent, preferably about 20 to 80, and more preferably about 30 to 70 weight percent of the composition.

A preferred stabilizer includes glycol stearate. The stabilizer, when used, is typically present in an amount from about 0.1 to 5 weight percent of the composition.

20

25

Preferred preservatives include tetrasodium ethylene-diamine tetraacetic acid (EDTA), methylparaben, benzophenone-4, methylchloroisothiazolinone, methylisothiazolinone, and the like, and mixtures thereof. Preservatives, when used, are typically present in an amount from about 0.01 to 6 weight percent, preferably about 0.05 to 4 weight percent, and more preferably from about 0.1 to 2 weight percent.

Preferred moisturizers include wheat protein (e.g., laurdimonium hydroxypropyl hydrolyzed wheat protein), hair keratin amino acids, sodium peroxylinecarbolic acid, panthenol, tocopherol (Vitamin E), dimethicone, and the like, and mixtures thereof. Sodium chloride may also be present, particularly when hair keratin amino acids are included as a moisturizer. Moisturizers are typically present in an amount from about 0.01 to 2 weight percent, preferably about 0.05 to 1.5 weight percent, more preferably from about 0.1 to 1 weight percent of the composition.

Preferred coloring agents include FD&C Green No. 3, Ext. D&C Violet No. 2, FD&C Yellow No. 5, FD&C Red No. 40, and the like, and mixtures thereof. The coloring agents are preferably used, and when used are typically present in an amount from about 0.001 to 0.1 weight percent, and preferably from about 0.005 to 0.05 weight percent of the composition.

Preferred anti-inflammatory agents include any pharmaceutically acceptable compounds suitable for administration orally or topically, preferably allantoin. The anti-inflammatory agents, when present, are used in an amount sufficient to inhibit or reduce inflammation, preferably in an amount from about 0.1 to 2 weight percent, preferably from about 0.3 to 1.5 weight percent, and more preferably from about 0.3 to 1 weight percent of the composition.

Anti-oxidants of both the enzymatic and non-enzymatic type may be 15 included in the compositions and methods of the invention. For example, superoxide dismutase (SOD), catalase, and glutathione peroxidase are natural enzymatic antioxidants used by the body that may be supplemented with the compositions herein. Suitable non-enzymatic anti-oxidants include such as Vitamin E (e.g., tocopherol), Vitamin C (ascorbic acid), carotenoids, Echinacoside and caffeoyl derivatives, oligomeric proanthocyanidins or proanthanols (e.g., grape seed extract), silymarin (e.g., milk thistle extract, Silybum marianum), ginkgo biloba, green tea polyphenols, and the like, and mixtures thereof. Carotenoids are powerful anti-oxidants, and they include beta-carotene, canthaxanthin, zeaxanthin, lycopen, lutein, crocetin, capsanthin, and the like. Indeed, any pharmaceutically acceptable compounds suitable for administration orally or topically may be used as an anti-oxidant in the compositions. Preferably, the anti-oxidant component includes Vitamin E, Vitamin C, or a carotenoid. The antioxidant component, when used, is present in an amount sufficient to inhibit or reduce 30 the effects of free-radicals at the scalp. The anti-oxidant component may be present in an amount from about 0.001 to 1 weight percent, preferably from about 0.01 to 0.5 weight percent of the composition.

Application of chemicals to the hair, such as for coloring or other hair treatment, requires the cuticles to be opened, which may be accomplished by using any of a variety of alkaline components suitable for application to the hair. It should be

understood that the application of the acidic component, however, will close or essentially close the cuticle incidental to the methods of managing scalp conditions, as discussed herein. Thus, chemical processing of the hair is rendered more difficult after application of the compositions of the present invention. However, when it is desired to specifically protect chemically processed hair, the compositions herein may be administered subsequent to, preferably immediately subsequent to, chemical processing of the hair. By "immediately" it is meant that the compositions herein are administered within 4 hours, preferably within 1 hour, and more preferably within 30 minutes of the chemical processing. Thus, in a preferred embodiment for protecting chemically processed hair, a patient may optionally have their hair conventionally shampooed and rinsed, chemically processed, and then have the compositions of the invention administered to protect the processing by closing or essentially closing the cuticle.

The term "scalp conditions" as used herein means conditions including dandruff, eczema, hair thinning, irritation, psoriasis, scaling, folliculitis, seborrhea or seborrheic dermatitis, and the like.

15

The terms "managing" or "management" as used herein includes one or more of the prevention, treatment, or modification of a condition.

The term "environmental damage" as used herein means weather or other damage to the scalp or hair from exposure to one or more of water, sun, pollution, overdrying, overteasing, and the like.

The term "therapeutically effective amount" means that amount of the pharmaceutical composition that provides a therapeutic benefit in the treatment, prevention, or management of scalp conditions.

The magnitude of a prophylactic or therapeutic dose of the composition in the acute or chronic management of scalp conditions will vary with the severity of the condition to be treated and the route of administration. The dose, and perhaps the dose frequency, will also vary according to the age, body weight, and response of the individual patient. In general, a preferred topical daily dose range, in single or divided doses, for the conditions described herein should be from about 1 mg to 20,000 mg, more preferably about 2,000 mg to 16,000 mg, and most preferably about 6,000 mg to 10,000 mg of the active components (i.e., excluding excipients and carriers).

Those of ordinary skill in the art will also understand that topical effectiveness of pharmaceuticals requires percutaneous absorption and bioavailability to the target site. Thus, the compositions and methods of the invention require penetration through the stratum corneum into the epidermal layers, as well as sufficient distribution to the sites targeted for pharmacologic action.

It is further recommended that children, patients aged over 65 years, and those with impaired renal or hepatic function initially receive low doses, and that they then be titrated based on individual response(s) or blood level(s). It may be necessary to use dosages outside these ranges in some cases, as will be apparent to those of ordinary skill in the art. Further, it is noted that the clinician or treating physician will know how and when to interrupt, adjust, or terminate therapy in conjunction with individual patient response.

The term "unit dose" is meant to describe a single dose, although a unit dose may be divided, if desired. About 1 to 2 unit doses of the present invention are typically administered per day, preferably about 1 dose per day.

Any suitable route of administration may be employed for providing the patient with an effective dosage of the composition according to the methods of the present invention, including oral, intraoral, rectal, parenteral, topical, epicutaneous, transdermal, subcutaneous, intramuscular, intranasal, sublingual, buccal, intradural, intraocular, intrarespiratory, or nasal inhalation and like forms of administration.

Topical administration is generally preferred for the compositions and methods of the invention, although oral administration is preferred for the 5-α reductase inhibitor. Suitable dosage forms include dispersions, suspensions, solutions, aerosols, sponges, cotton applicators, and the like, with topical dosage forms such as shampoos being preferred. A sunscreen dosage form is one preferred embodiment for methods and compositions for the management of thinning hair, since extra protection is required when more scalp is exposed to harmful ultraviolet radiation. In a sunscreen dosage form, it is preferred to include benzophenone-4, which provides the scalp with some sunscreen protection factor from ultraviolet radiation.

The pharmaceutical compositions used in the methods of the present invention include the active ingredients described above, and may also contain

35

pharmaceutically acceptable carriers, excipients and the like, and optionally, other therapeutic ingredients.

The term "pharmaceutically acceptable salt" refers to a salt prepared from

pharmaceutically acceptable non-toxic acids or bases including inorganic or organic

acids. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic,

sulfuric, and phosphoric. Appropriate organic acids may be selected, for example, from
aliphatic, aromatic, carboxylic and sulfonic classes of organic acids, examples of which
are formic, acetic, propionic, succinic, glycolic, glucoronic, maleic, furoic, glutamic,
benzoic, anthranilic, salicylic, phenylacetic, mandelic, embonic (pamoic),
methanesulfonic, ethanesulfonic, pantothenic, benzenesulfonic, stearic, sulfanilic,
algenic, and galacturonic. Examples of such inorganic bases, for potential salt
formation with the sulfate or phosphate compounds of the invention, include metallic
salts made from aluminum, calcium, lithium, magnesium, potassium, sodium, and zinc.
Appropriate organic bases may be selected, for example, from N,Ndibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine,
meglumaine (N-methylglucamine), and procaine.

The compositions for use in the methods of the present invention may include components such as suspensions, solutions and elixirs; aerosols; or other suitable carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents, and the like, with the topical preparations being preferred.

Because of its ease of administration, shampoo represents the most advantageous topical dosage unit form, in which case liquid pharmaceutical carriers may be employed in the composition. These shampoos may be prepared as rinse-off or leave-on after-shampoo products, as well as two stage treatment shampoos.

30 Additionally, the composition may also be prepared as a stand-alone conditioner or preor post-shampoo conditioner. In a preferred embodiment, the compositions are administered as a shampoo, then as a conditioner after the shampoo is rinsed away, and optionally in a higher concentration form, such as a gel, subsequently thereto. Each of these forms is well understood by those of ordinary skill in the art, such that dosages may easily be prepared to incorporate the pharmaceutical composition of the invention.

Pharmaceutical compositions for use in the methods of the present invention suitable for topical administration may be presented as discrete units including aerosol sprays, each containing a predetermined amount of the active ingredient, as a powder, stick, or granules, as creams (e.g., a conditioner), pastes, gels, lotions (e.g., a sunscreen), syrups, or ointments, on sponges or cotton applicators, or as a solution or a suspension in an aqueous liquid, a non-aqueous liquid, an oil-in-water emulsion, or a water-in-oil liquid emulsion. Such compositions may be prepared by any of the methods of pharmacy, but all methods include the step of bringing into association the carrier(s) with the active ingredient, which constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired presentation.

Other suitable dosage forms include tablets, troches, capsules, patches, gel caps, magmas, lozenges, plasters, discs, suppositories, nasal or oral sprays, and the like. Tablets, capsules, and gel caps are the preferred oral dosage unit form for the  $5-\alpha$  reductase inhibitor, in which case solid pharmaceutical carriers may be employed. If desired, tablets may be coated by standard aqueous or nonaqueous techniques.

In addition to the common dosage forms set out above, the compound for use in the methods of the present invention may also be administered by controlled release means and/or delivery devices such as those described in U.S. Patent Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, the disclosures of which are expressly incorporated herein by reference thereto.

For example, a tablet may be prepared by compression or molding, optionally, with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, surface active or dispersing agent. Molded tablets may be made by molding, in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent.

35

15

25

Desirably, each unit dose, e.g., shampoo, contains from about 1 mg to 2,000 mg of the active ingredient, preferably about 200 mg to 1,600 mg, and more preferably about 600 mg to 1,000 mg of the composition.

5

## **EXAMPLES**

The invention is further defined by reference to the following examples describing in detail the preparation of the compound and the compositions used in the methods of the present invention, as well as their utility. The examples are representative, and they should not be construed to limit the scope of the invention.

Example 1: Shampoo for Dandruff Treatment

A pharmaceutical composition according to the invention may be prepared 15 in the form of a shampoo formulated for dandruff management as set forth below:

		Ingredient	Trade Name/Supplier	% by Weight
	Part A	Deionized Water	N/A	47
20		Guar Hydroxypropyltrimonium Chloride	JAGUAR C-13S	0.1
		Glycolic Acid	GLYPURE-70% Glycolic Acid	1.5
		Methylparaben	Methylparaben	0.2
		Sodium Laureth Sulfate	CARSONOL SLES-2	28
25		Glycol Distearate	LEXEMUL EGDS	1
		Cocamidopropyl Betaine	LONZAINE C	5
	Part B	Lauramide Diethanolamine (DEA)	MONAMID 716	2
	Part C	Salicylic Acid	Salicylic Acid, powdered, USP	1.2
30	•	Sodium Laureth Sulfate	CARSONOL SLES-2	10
	Part D	Allantoin	ALLANTOIN	0.1
		Panthenol	DL-PANTHENOL 50%	0.3
35		Dimethicone (and) Laureth-4 and Laureth-23	DOW CORNING 1664	0.05
	Part E	Clotrimazole	Clotrimazole, USP	0.5
		Alcohol (denatured)	SD ALCOHOL 40, Anhydrous	1.5

Part F	Glycolic Acid	GLYPURE-70% Glycolic Acid	0.15
	Phenoxyethanol	Phenoxyethanol	0.7
	Methylchloroisothiazolinone and Methylisothiazolinone	KATHON CG	0.05
5	Retinyl Palmitate	Vitamin A Palmitate, type P1.7	0.02
	Tocopherol	Vitamin E USP, #60525	0.02
	Benzophenone-4	UVINUL MS-40	0.1
•	FD&C Green No. 3 (CI 42053)	FD&C Green No. 3 (1.0% Solution)	0.02
0	Ext. D&C Violet No. 2 (CI 60730)	Ext. D&C Violet No. 2 (0.10% Solution)	0.2
	Fragrance (Parfum)	Natural Fragrance - Haarmann & Reimer #A40758/778926	0.3
		•	100%

15 JAGUAR C-13S is commercially available from Rhone-Poulenc N. American Chemicals of Cranbury, NJ; GLYPURE is commercially available from DuPont of Wilmington, DE; CARSONOL SLES-2 and LONZAINE C are commercially available from Lonza, Inc. of Fair Lawn, NJ; LEXEMUL EGDS is commercially available from Inolex Chemical Co. of Philadelphia, PA; MONAMID 716 is commercially available from Mona Industries, Inc. of Paterson, NJ; Allantoin is commercially available from ISP Corp. of Bound Brook, NJ; DL-PANTHENOL and UVINUL MS-40 are commercially available from BASF Corp. of Budd

20 Lake, NJ; DOW CORNING 1664 is commercially available from Dow Corning Corp. of Auburn, MI; KATHON CG is commercially available from Rohm & Haas Co. of Philadelphia, PA; FD&C Green No. 3 and Ext. D&C Violet No. 2 are commercially available from Hilton-Davis Co. of Cincinnati, OH; and Natural Fragrance #A4-758/778926 is commercially available from Haarmann & Reimer of Holzminden, Germany (subsidiary of Bayer Aktiengesellschaft).

Deionized water may be metered into the processing tank and high speed mixing subsequently begun. Guar hydroxypropyltrimonium and glycolic acid are added one at a time. Once the guar component is completely dispersed, the mixture is heated to 75°C. The remaining Part A ingredients are then added and mixed until all the solids are dissolved. The mixture is cooled to 60°C and the Part B ingredients are added and mixed until uniform. The mixture is then cooled to 50°C. In a separate vessel, Part C is premixed until homogeneous, then added to the mixture of Parts A & B. Parts A, B, and C are mixed until uniform. The Part D ingredients are added and mixed until uniform, then cooled to 40°C. Premixed Part E is added and mixed until uniform, the Part F ingredients are added and mixed, and the composition is then cooled to 35°C. The appearance should be an aqua blue, opaque, pearlized, semiviscous liquid having a pH at 25°C of between 4 to 5 and a viscosity between 5,000 to 9,000 cps (RVT: #4 @ 10 rpm @ 25°C).

Example 2: Shampoo for Thinning Hair

A pharmaceutical composition according to the invention may be prepared in the form of a shampoo formulated for management of thinning hair as set forth below:

5				
		Ingredient	Trade Name/Supplier	% by Weight
	Part A	Deionized water	N/A	51.9
		Polyquaternium-10	UCARE Polymer JR-125	0.15
0		Glycolic Acid	GLYPURE-70% Glycolic Acid	1
		Methylparaben	N/A	0.2
		Sodium Laureth Sulfate	CARSONOL SLES-2	35
	•	Glycol Stearate	LEXEMUL EGMS	0.75
		Phytantriol	N/A	0.1
5		Sodium Lauroyl Sarcosinate	HAMPOSYL L-30	1.5
		Cocamidopropyl Betaine	LONZAINE C	5
		Panthenyl Ethyl Ether	Ethyl Panthenol	0.2
0		Tetrasodium Ethylene-Diamine- Tetraacetic Acid (EDTA)	HAMP-ENE 220	0.03
U		Benzophenone-4	Uvinul MS-40	0.1
	Part B	Lauramide Diethanolamine	MONAMID 716	2.2
	Part C	Lactamide Monoethanolamine (MEA)	PARAPEL LAM-100	0.6
5		Sodium Peroxylinecarbolic Acid (PCA)	AJIDEW N-50	0.4
<del></del>		Hair Keratin Amino Acids and Sodium Chloride	CROTEIN HKP	0.1
	Part D	Saw Palmetto (Serenoa Serrulata) Extract	Saw Palmetto Berry Glycolic Extract	0.1
0		Methylchloroisothiazolinone and Methylisothiazolinone	KATHON CG	0.05
		FD&C Green No. 3 (CI 42053)	FD&C Green No. 3 (1.0% Solution)	0.005
		FD&C Yellow No. 5 (CI 19140)	FD&C Yellow No. 5 (1.0% Solution)	0.007
5	Part E	Menthol	Menthol Crystals, USP	0.3

PCT/US99/16866 WO 00/06144

Essential Oil Blend #EE-50098 Cinnamon (Cinnamomum Cassia) Oil, 0.5 Grapefruit (Citrus Grandis) Oil, Orange (Citrus Aurantium Dulcis) Oil. Petitgrain (Citrus Aurantium Amara) Oil, Peppermint (Mentha Piperita) Oil, Eucalyptus Globulus Oil, Clove (Eugenia Caryophyllus) Oil, and Rosemary (Rosmarinus Officinalis) Oil Fragrance - Haarmann & 0.3 Fragrance (Parfum)

Reimer #A40758/778926

10

5

100%

UCARE Polymer JR-125 is commercially available from Amerchol Corp. of Edison, NJ; PHYTRANTIOL and ETHYL PANTHENOL are commercially available from Roche Vitamins Inc. of Parsippany, NJ; HAMPOSYL L-30 is commercially available from Hampshire of Lexington, MA: HAMP-ENE 220 is commercially available from Akzo Nobel Inc. of Dobbs Ferry, NY; PARAPEL LAM-100 is commercially available from Bernel Chemical Co. of Englewood, NJ; AJIDEW N-50 is commercially available from Ajinomoto USA Inc. of Teaneck, NJ; CROTEIN HKP is commercially available from Croda Inc. of Parsippany, NJ; Saw Palmetto is commercially available from Chart at Patterson, NJ; and FD&C Green No. 3 and Yellow No. 5 are commercially available from Hilton Davis Co. of Cincinnati, OH.

Deionized water may be metered into the processing tank and high speed mixing subsequently begun. Polyquaternium-10 and glycolic acid are added one at a time. Once the polymer is completely dispersed, the mixture is heated to 75°C. The remaining Part A ingredients are then added and mixed until all the solids are dissolved. The mixture is cooled to 60°C and the Part B ingredients are added and mixed until uniform. The mixture is then cooled to 50°C. In a separate vessel, Part C is premixed until homogeneous, then added to the mixture of Parts A & B. Parts A, B, and C are mixed until uniform. The Part D ingredients are added and mixed until uniform, then cooled to 40°C. Premixed Part E is added and mixed until uniform, the Part F ingredients are added and mixed, and the composition is then cooled to 35°C. The appearance should be a light green, opaque, pearlized, semi-viscous liquid having a 30 pH at 25°C of between 5.4 to 6.1 and a viscosity between 6,000 to 12,000 cps (RVT: #5 @ 10 rpm @ 25°C).

# Example 3: Shampoo Formulation for Chemically Treated Hair

A pharmaceutical composition according to the invention may be formulated as a shampoo for protecting chemically treated hair as set forth below:

		Ingredient	Trade Name/Supplier	<u>% by</u> <u>Weight</u>
	Part A	Deionized water	N/A	86.4%
_	•	Polyquaternium-10	UCARE Polymer JR-125	0.2
5		Glycolic Acid	GLYPURE-70% Glycolic Acid	0.1
		Tetrasodium Ethylene-Diamine- Tetraacetic Acid (EDTA)	HAMP-ENE 220	0.01
		Sodium Laureth Sulfate	STEOL CS-230 PCK	10
_		Glycol Stearate	LEXEMUL EGMS	0.1
0		Methylparaben	N/A	0.05
		Sodium Lauroyl Sarcosinate	HAMPOSYL L-30	0.1
		Panthenol	Liquid DL-Panthenol 50%	0.5
		Benzophenone-4	UVINUL MS-40	0.05
5	Part B	Disodium Cocoamphodiacetate	MIRANOL C2M CONC. NP	1
	Part C	Lactamide Monoethanolamine (MEA)	PARAPEL LAM-100	0.5
		Sodium Peroxylinecarbolic Acid (PCA)	AJIDEW N-50	0.5
		Hair Keratin Amino Acids and Sodium Chloride	CROTEIN HKP	0.05
0		Laudimonium Hydroxypropyl Hydrolyzed Wheat Protein	HYDROTRITICUM QL	0.01
	Part D	Methylchloroisothiazolinone and Methylisothiazolinone	KATHON CG	0.05
		FD&C Red No. 40 (CI 16035)	FD&C Red No. 40 (1.0% Solution)	0.02
5	Part E	Jasmine (Jasminum Officinale) Oil, Palmarosa (Cymbopogon Martini) Oil, Sandalwood (Santalum Album) Oil, Patchouli (Pogostemon Cablin) Oil, Nutmeg (Myristica Fragrans) Oil, Grapefruit (Citrus Grandis) Oil, Orris (Iris Florentina) Root Oil, Carnation	Essential Oil Blend "Passion"	0.3
0		(Dianthus Caryophyllus) Oil		
		Phytantriol	N/A	0.02
	Part F	Glycolic acid	GLYPURE-70% glycolic acid	0.1
	Part G	Deionized water	N/A	0.1
5		Sodium chloride	N/A	0.1
-				100%

HYDROTRITICUM QL is commercially available from Croda Inc. of Parsippany, NJ; MIRANOL C2M is commercially available from Rhone-Poulenc N. American Chemicals of Cranbury, NJ; Phytantriol is commercially available from Roche Vitamins Inc. of Parsippany, NJ; and STEOL CS-230 PCK is commercially available from Stepan Co. of Northfield, IL.

5

Deionized water may be metered into the processing tank and high speed mixing subsequently begun. Polyquaternium-10 and glycolic acid are added one at a time. Once the polymer is completely dispersed, the mixture is heated to 75°C. The remaining Part A ingredients are then added and mixed until all the solids are dissolved. The mixture is cooled to 60°C and the Part B ingredients are added and mixed until uniform. The mixture is then cooled to 50°C. Part C is then added to the mixture of Parts A & B and mixed until uniform. The Part D ingredients are added and mixed until uniform, then cooled to 40°C. Premixed Part E is added and mixed until uniform, the Part F ingredients are added in increments as need to obtain the desired pH of 5.7 to 6.5 and mixed until uniform. Then Part G is added in increments to obtain the desired viscosity of 8,000 to 12,000 cps (RVT#5 @ 10 rpm @ 25°C), and the composition is then mixed and cooled to 35°C. The appearance should be a peach-colored, opaque, pearlized, semi-viscous liquid having a pH at 25°C of between 5.7 to 6.5 and a viscosity between 8,000 to 12,000 cps (RVT: #5 @ 10 rpm @ 25°C).

Example 4: Professional Scalp Formulation for Dandruff

A pharmaceutical composition according to the invention may be formulated in a more potent concentration as a shampoo for professional management of dandruff as set forth below:

	Ingredient		Trade Name/Supplier	% By Weight
30	Part A	Deionized water	N/A	74.6%
		Hydroxyethylcellulose	CELLOSIZE QP52,000H	0.3
35	Part B	Disodium Ethylene-Diamine- Tetraacetic Acid (EDTA)	HAMP-ENE Na₂	0.03
33		Glycerin	Glycerine 99.5%	1

		Glycolic Acid	GLYPURE-70% Glycolic Acid	10
	Part C	Deionized water	N/A	1 .
5		Sodium Hydroxide	Sodium Hydroxide, pellets (USP/NF)	1
	Part D	Alcohol (denatured)	SD Alochol 40-B, Anhydrous	10
10		Salicylic Acid	Salicylic Acid, powder (USP/NF)	1.8
	Part E	PPG-5-Ceteth-20	PROCETYL AWS	0.2
		Lemongrass (Cymbopogon Schoenanthus) Oil	Lemongrass Oil #65161	0.02
15		Peppermint (Mentha Piperita) Oil	Peppermint Oil #097973- 0594N	0.04
		Geranium Maculatum Oil	Geranium Oil #01491	0.02
				100%

HAMP-ENE Na<sub>2</sub> is commercially available from Akzo Nobel Inc. of Dobbs Ferry, NY; and PROCETYL AWS is commercially available from Croda Inc. of Parsippany, NJ.

Deionized water may be metered into the processing tank and high speed mixing subsequently begun. CELLOSIZE QP52,000H is added and heated at 70°C until the solids are dissolved. The mixture is cooled to 40°C. The Part B ingredients are added and mixed until uniform. Premixed Part C is then added in increments to obtain the desired pH below, and mixed until uniform. The mixture is cooled to 25°C and premixed part D is slowly added and mixed until uniform. Premixed Part E is added and mixed until uniform. The composition should be colorless, clear, semi-viscous liquid having a pH at 25°C between 3.4 to 3.8 and a viscosity between 1,500 to 2,5000 cps (RVT#5 @ 10 rpm @ 25°C).

# Example 5: Scalp Formulation for Thinning Hair

A pharmaceutical composition according to the invention may be formulated as a shampoo for management of thinning hair as set forth below:

		Ingredient	Trade Name	% By Weight
	Part A	Deionized water	N/A	82.3%
5	Part B	Propylene Glycol	N/A	5
5		Methylparaben	N/A	0.2
	Part C	Menthol	Menthol Crystals, USP	0.03
		Ethyl Nicotinate		0.05
LO		Cinnamon (Cinnamomum Cassia) Oil, Grapefruit (Citrus Grandis) Oil, Orange (Citrus Aurantium Dulcis) Oil, Petitgrain (Citrus Aurantium Amara) Oil	Essential Oil Blend #EE-50098	0.04
		Polysorbate 80	TWEEN 80	3
15		Phytantriol	Phytantriol	0.2
		Biotin	N/A	0.1
		Salicylic Acid	Salicylic Acid, powder, USP/NF	0.5
		Capsicum Frutescens Extract and Safflower (Carthamus Tinctorius) Oil	Actiphyte of Capsicum, Lipo S	0.05
0	Part D	Allantoin	Allantoin	0.5
		Tetrasodium EDTA	HAMP-ENE 220	0.2
		Lactamide Monoethanolamine	PARAPEL LAM-100	1.5
		Panthenyl Ethyl Ether	Ethyl Panthenol	1.5
5		Diazolidinyl Urea	GERMALL II	0.3
		Cocamidopropyl PG-Dimonium Chloride Phosphate	PHOSPHOLIPID PTC	1.08
		Saw Palmetto (Sereno Serrulata) Extract	Saw Palmetto Extract	0.5
		Cyanocobalamin	Vitamin B <sub>12</sub>	0.02
0		Yeast Extract (Faex)	NAYAD S	1
-		Yeast Extract (Faex)	RESPIROGEN	2
				100%

PHOSPHOLIPID PTC is commercially available from Mona Industries Inc. of Patterson, NJ; and RESPIROGEN is commercially available from Immudyne, Inc. of Houston, TX.

Preparation Procedure:

Meter deionized water into the processing tank. Start mixing. In a separate vessel, heat Part B to 60°C until all the solids are dissolved. Slowly add Part B to Part A with high speed mixing. Mix until uniform. In a separate vessel, premix Menthol, Ethyl Nicotinate and the Essential Oil Blend until all the solids are dissolved. Add the remaining Part C ingredients. Mix until uniform. Slowly add Part C to the batch. Mix until clear and homogeneous. Add Part B ingredients in the given order, mixing well after each addition. Mix until completely uniform. The resulting composition should be a light golden yellow, clear, non-viscous liquid having a pH at 25°C of between 3.6 to 4.2.

# Example 6: Conditioner Formulation for Thinning Hair

A pharmaceutical composition according to the invention may be formulated as a conditioner for management of thinning hair as set forth below:

15

20

25

30

35

	-	Ingredient	Trade Name	% By Weight
	Part A	Deionized water	N/A	79.5 %
		Hydroxyethylcellulose	CELLOSIZE POLYMER PCG-10	1
		Methylparaben	N/A	0.2
5		Benzophenone-4	UVINUL MS-40	0.1
		Glyceryl Stearate	RITA GMS-55G	2
		Stearalkonium Chloride	MAQUAT SC-18 85%	2
		Cetyl Alcohol	LANETTE 16	2.5
		Stearyl Alcohol	LANETTE 18	2.5
		Glycerin	Glycerine 99.5%	1
)	Part B	Laurdimonium Hydroxypropyl Hydrolyzed Wheat Protein	HYDROTRITICUM QL	. 1
		Panthenol	DL-PANTHENOL 50%	1
		Hydrolyzed Soy Protein	HYDROSOY 2000 SF	1
		Hydrolyzed Wheat Protein Hydroxypropyl Polysiloxane	CRODASONE W	1
		Hair Keratin Amino Acids and Sodium Chloride	CROTEIN HKP Powder	1
;	Part C	Methylchloroisothiazolinone and Methylisothiazolinone	KATHON CG	0.05
		Benzalkonium Chloride	BTC-50	0.5
		Biotin	Biotin	0.05
		Ethyl Nicotinate	Ethyl Nicotinate	0.05
		Witch Hazel (Hamamelis Virginiana) Distillate	Witch Hazel Distillate E02	2
		FD&C Blue No. 1 (CI 42090)	FD&C Blue No. 1	0.006
)			(1.0% Solution)	
		FD&C Yellow No. 5 (CI 19140)	FD&C Yellow No. 5	0.006
	Part D	Cinnamon (Cinnamomum Cassia) Oil,	(1.0% Solution)	
		Grapefruit (Citrus Grandis) Oil, Orange	Essential Oil Blend	0.1
		(Citrus Aurantium Dulcis) Oil, Petitgrain	#EE-50098	
		(Citrus Aurantium Amara) Oil, Peppermint		
		(Mentha Piperita) Oil, Eucalyptus Globulus		
		Oil, Clove (Eugenia Caryophyllus) Oil,		
,		Rosemary (Rosmarinus Officinalis) Oil Menthol		
		Fragrance (Parfum)	Menthol Crystals, USP	0.5
	Part E	Glycolic Acid	Fragrance - HAARMANN & REIMER #A40758/778926	0.5
			GLYPURE-70% Glycolic Acid	0.5
				100%

CRODASONE W is commercially available from Croda Inc. of Parsippany, NJ.

# Preparation Procedure:

Meter deionized water into the processing tank. Start high speed mixing. Sprinkle-in CELLOSIZE Polymer PCG-10. When the Polymer is completely dispersed, heat to 75°C. Add the remaining Part A ingredients. Mix at 75°C for 30 minutes until homogeneous. Cool at 50°C. Add Part B ingredients. Mix until uniform. Cool to 40°C. Add Part C ingredients. Mix until uniform. In a separate vessel, premix Part D until all the solids are dissolved. Add to the batch. Mix until uniform. Add Part E in increments

as needed to obtain the desired pH. Continue mixing and cooling to 35°C. The resulting composition should have a very light green, opaque, semi-viscous appearance having a pH at 25°C of between 3.8 to 4.5 and a viscosity between 7,000 to 11,000 cPs (RVT: #4 @ 10 rpm @ 25°C).

# 5 Example 7: Professional Scalp Treatment for Chemically Treated Hair

A pharmaceutical composition according to the invention may be formulated in a higher concentration of acid as a professional treatment for protection of chemically treated hair as set forth below:

10				<del></del>
		Ingredient	Trade Name	% By Weight
	Part A	Deionized water	N/A	73%
		Hydroxyethylcellulose	CELLOSIZE QP52,000H	1
15	Part B	Disodium EDTA	HAMP-ENE Na <sub>2</sub>	0.2
		Glycerin	Glycerine 99.5%	5
		Panthenol	Liquid DL-Panthenol 50%	1
		Quaternium-75	FINQUAT CT	1.5
20		Cetrimonium Chloride	CARSOQUAT CT-429	1.5
		Glycolic Acid	GLYPURE-70% Glycolic Acid	12
	Part C	Deionized water	N/A	2
25		Sodium Hydroxide	Sodium Hydroxide, pellets, USP/NF	1.5
	Part D	PPG-5-Ceteth-20	PROCETYL AWS	1
		Phytantriol		0.2
		Ylang Ylang (Cananga Odorata) Oil	Ylang Ylang Oil #65040	0.04
30		Jasmine (Jasminum Officinale) Oil	Jasmine Oil #65072	0.01
		Chamomile (Anthemis Nobilis) Oil	Chamomile Oil #112528- 0395N	0.02
35				100%

FINQUAT CT is commercially available from Finetex Inc. of Elmwood Park, NJ.

## Preparation Procedure:

Meter deionized water into the processing tank. Start high speed mixing. Sprinkle-in CELLOSIZE QP52,000H. Heat to 70°C. Mix until all the solids are completely dissolved, and the batch is clear and uniform. Cool to 40°C. Add Part B ingredients. Mix until uniform. Slowly add premixed Part C ingredients in increments as needed to obtain the desired pH. Mix until uniform. Add premixed Part D. Continue mixing and cooling to 35°C. The resulting composition should be a colorless to pale straw-colored, clear, semi-viscous liquid having a pH at 25°C of between 3.4 to 3.8 and a viscosity between 2,500 to 3,500 cPs (RVT: #3 @ 10 rpm @ 25°C).

10

# Example 8: Professional Scalp Management of Thinning Hair

A pharmaceutical composition according to the invention may be formulated in a higher concentration of acid for professional management of thinning hair as set forth below:

7	E
_	2

		Ingredient	Trade Name	% By Weight
	Part A	Deionized water	N/A	77.5%
20		Hydroxyethylcellulose	CELLOSIZE QP52,000H	1
	Part B	Disodium EDTA	HAMP-ENE Na <sub>2</sub>	0.2
		Glycerin	Glycerine 99.5%	5
		Panthenyl Ethyl Ether	Ethyl Panthenol	1
25		Glycolic Acid	GLYPURE-70% Glycolic Acid	12
	Part C	Deionized water	N/A	1.8
		Sodium Hydroxide	Sodium Hydroxide, pellets, USP/NF	1
30	Part D	PPG-5-Ceteth-20	PROCETYL AWS	0.2
		Phytantriol	Phytantriol	0.2
		Ethyl Nicotinate	Ethyl Nicotinate	0.05
		Menthol	Menthol Crystals, USP	0.03
35		Cinnamon (Cinnamomum Cassia) Oil	Cinnamon Oil #65028	0.02
		Rosemary (Rosmarinus Officinalis) Oil	Rosemary Oil #65138	0.03

#097973-0594N	
	100%

Preparation Procedure:

Meter deionized water into the processing tank. Start high speed mixing. Sprinkle-in Cellosize QP52,000H. Heat to 70°C. Mix until all the solids are completely dissolved, and the batch is clear and uniform. Cool to 40°C. Add Part B ingredients. Mix until uniform. Slowly add premixed Part C in increments as needed to obtain the desired pH.

10 Mix until uniform. Add premixed Part D. Continue mixing and cooling to 35°C. The resulting composition should be a colorless to pale straw-colored, clear, semi-viscous liquid having a pH at 25°C between 3.4 to 3.8 and a viscosity between 2,200 to 3,500 cPs (RVT: #3 @ 10 rpm @ 25°C).

# Example 9: Conditioner for Chemically Treated Hair

A pharmaceutical composition according to the invention may be formulated as a conditioner for chemically treated hair as set forth below:

20		Ingredient	Trade Name	% By Weight
	Part A	Deionized water	N/A	88%
		Guar Hydroxypropyltrimonium Chloride	JAGUAR C-13S	0.5
25		Glycolic Acid	GLYPURE-70% Glycolic Acid	0.15
		Methylparaben	N/A	0.15
	•	Stearalkonium Chloride	MAQUAT SC-18 85%	2
		Cetyl Alcohol	LANETTE 16	2
30	÷	Stearyl Alcohol	LANETTE 18	2
50		Trimethylsilylamodimethicone	SF 1708-D1	0.5
		Sodium PCA	AJIDEW N-50	0.05
		Benzophenone-4	UVINUL MS-40	0.1
35	Part B	Hydrolyzed Wheat Protein and Hydrolyzed Wheat Starch	CROPEFTIDE W	0.1

100%

		Hydrolyzed Wheat Protein Hydroxypropyl Polysiloxane	CRODASONE W	0.5
5	٠	Panthenol	Liquid DL-Panthenol 50%	1
		Hydrolyzed Soy Protein	HYDROSOY 2000 SF	1
		Phytantriol	Phytantriol	0.3
		Polyquaternium-11	GAFQUAT 755N	1
10	Part C	Methylchloroisothiazolinone and Methylisothiazolinone	KATHON CG	0.05
		Benzalkonium Chloride	BTC-50	0.3
		FD&C Red No. 40 (Cl 16035)	FD&C Red No. 40 (1.0% Solution)	0.012
20		Jasmine (Jasminum Officinale) Oil, Palmarosa (Cymbopogon Martini) Oil, Sandalwood (Santalum Album) Oil, Patchouli (Pogostemon Cablin) Oil, Nutmeg (Myristica Fragrans) Oil, Grapefruit (Citrus Grandis) Oil, Orris (Iris Florentina) Root Oil, Carnation (Dianthus Caryophyllus) Oil	Essential Oil Blend "Passion"	0.3

GAFQUAT 775N is commercially available from ISP Corp. of Bound Brook, NJ.

## Preparation Procedure:

Meter deionized water into the processing tank. Start high speed mixing. Sprinkle-in Jaguar C-13S. Add Glypure. When the Jaguar is completely dispersed, heat to 75°C. Add the remaining Part A ingredients. Mix at 75°C for 30 minutes until homogeneous. Cool to 50°C. Add Part B ingredients. Mix until uniform. Cool to 40°C. Add Part C ingredients. Continue mixing and cooling to 35°C. The resulting composition should be light peach-colored, opaque, semi-viscous, having a pH at 25°C between 4 to 4.8 and a viscosity between 8,000 to 12,000 cPs (RVT: #4 @ 10 rpm @ 25°C).

35

25

Various modifications of the invention in addition to those shown and described herein will be apparent to those skilled in the art from the foregoing description. Such

modifications are also intended to fall within the scope of the appended claims. The foregoing disclosure includes all the information deemed essential to enable those skilled in the art to practice the claimed invention.

## THE CLAIMS

What is claimed is:

1. An anti-hair thinning pharmaceutical composition for administration to a scalp comprising:

an acidic component comprising a hydroxy acid or tannic acid, or a pharmaceutically acceptable salt thereof, in an amount sufficient to exfoliate at least a portion of the scalp;

a niacin component present in an amount sufficient to locally increase blood circulation; and

a  $5-\alpha$  reductase inhibitor in an amount sufficient to inhibit conversion of testosterone to dihydro-testosterone.

- 2. The pharmaceutical composition of claim 1, wherein the composition further comprises a pharmaceutically acceptable carrier or excipient.
  - The pharmaceutical composition of claim 1, wherein the acidic component comprises an alpha-hydroxy acid, beta-hydroxy acid, or tannic acid, the niacin component comprises nicotinate, and the 5-α reductase inhibitor comprises at least one of finasteride or
     Saw Palmetto Extract.
    - 4. The pharmaceutical composition of claim 3, wherein the acidic component comprises glycolic acid, lactic acid, citric acid, salicylic acid, or tannic acid.
- 5. The pharmaceutical composition of claim 1, wherein the acidic component is present in an amount from about 0.1 to 8 weight percent, the niacin component is present in an amount from about 0.01 to 1 weight percent, and the 5-α reductase inhibitor is present in an amount from about 0.1 to 1 weight percent, of the composition.
- The pharmaceutical composition of claim 1, further comprising at least one of a surfactant, a stabilizer, a preservative, a moisturizer, anti-inflammatory agent, anti-

oxidant, and a coloring agent, which together may be present in an amount from about 10.1 to 99.1 weight percent of the composition.

- 7. The pharmaceutical composition of claim 1, wherein the acidic component
   5 and the niacin component are adapted for topical administration and the 5-α reductase inhibitor is adapted for oral administration.
  - 8. A shampoo, cream, or gel comprising the pharmaceutical composition of claim 1.

10

- 9. A method of managing a scalp condition which comprises administering to a patient a therapeutically effective amount of an acidic component comprising a hydroxy acid or tannic acid, or a pharmaceutically acceptable salt thereof, a vitamin A component, and an anti-growth agent to inhibit fungus, yeast, bacteria, or a mixture thereof that may be present adjacent the scalp.
- 10. The method of claim 9, wherein the scalp condition treated is at least one of dandruff, seborrheic dermatitis, psoriasis, or folliculitis.
- 20 11. The method of claim 9, wherein the administering is topical.
  - 12. The method of claim 11, wherein the administering is by at least one of a shampoo, aerosol spray, gel, paste, cream, lotion, sponge, emulsion, or ointment.
- 25 13. The method of claim 9, wherein about 1 mg to 10,000 mg of the acidic component, vitamin A component, and anti-growth agent are together administered.
  - 14. The method of claim 9, wherein the acidic component, vitamin A component, and anti-growth agent are administered concurrently.

30

15. The method of claim 9, wherein the acidic component, vitamin A component, and anti-growth agent are administered concurrently with at least one additional pharmaceutical composition for the prevention or treatment of a scalp condition.

- 5 16. The method of claim 9, which further comprises administering at least one of a surfactant, stabilizer, preservative, moisturizer, anti-inflammatory agent, anti-oxidant, or coloring agent.
- 17. The method of claim 9, wherein the acidic component comprises an alpha-10 hydroxy acid, beta-hydroxy acid, or tannic acid, the vitamin A component comprises retinyl palmitate, and the anti-growth agent comprises at least one of triclosan or clotrimazole.
- 18. A method of managing hair thinning which comprises administering to a patient a therapeutically effective amount of an acidic component comprising a hydroxy acid or tannic acid, or a pharmaceutically acceptable salt thereof, a niacin component present in an amount sufficient to locally increase blood supply, and a 5-α reductase inhibitor.
  - 19. The method of claim 18, wherein the administering is topical.

20

20. The method of claim 18, wherein the acidic component and niacin component are topically administered and the 5-α reductase inhibitor is orally administered.

- The method of claim 19, wherein the administering is by at least one of a shampoo, aerosol spray, gel, paste, cream, lotion, sponge, emulsion, or ointment.
  - 22. The method of claim 18, wherein about 1 mg to 10,000 mg of the acidic component, niacin component, and 5-α reductase inhibitor are together administered.
- 30 23. The method of claim 18, wherein the acidic component, niacin component, and 5-α reductase inhibitor are administered concurrently.

24. The method of claim 18, wherein the acidic component, niacin component, and 5-α reductase inhibitor are administered concurrently with at least one additional pharmaceutical composition for the prevention or treatment of a scalp condition.

- 5 25. The method of claim 18, which further comprises administering at least one of a surfactant, stabilizer, preservative, moisturizer, anti-inflammatory agent, anti-oxidant, or coloring agent.
- 26. The method of claim 18, wherein the acidic component comprises an alphahydroxy acid, beta-hydroxy acid, or tannic acid, the niacin component comprises nicotinate, and the 5-α reductase inhibitor comprises at least one of finasteride or Saw Palmetto Extract.
- 27. A method of treating chemically processed hair having a plurality of cuticle which comprises administering to a patient an amount of an acidic component comprising a hydroxy acid or tannic acid, or a pharmaceutically acceptable salt thereof, in an amount sufficient to essentially close the cuticle and inhibit modification of the chemically processed hair.
- 28. The method of claim 27, wherein the administering is topical.
  - 29. The method of claim 27, wherein the administering is by at least one of a shampoo, aerosol spray, gel, paste, cream, lotion, sponge, emulsion, or ointment.
- The method of claim 27, wherein about 1 mg to 10,000 mg of the acidic component is administered.
- 31. The method of claim 27, wherein the acidic component is administered concurrently with at least one additional pharmaceutical composition for the prevention or treatment of a scalp condition.

32. The method of claim 27, which further comprises administering at least one of a surfactant, stabilizer, preservative, moisturizer, anti-inflammatory agent, anti-oxidant, or coloring agent.

5 33. The method of claim 27, wherein the acidic component comprises an alphahydroxy acid, beta-hydroxy acid, or tannic acid.

10

15

20

25

30

## INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/16866

· · · · · · · · · · · · · · · · · · ·			
	ASSIFICATION OF SUBJECT MATTER		
	: A61K 31/07, 31/19, 31/35, 31/44, 31/075 :514/284, 356, 460, 568, 574, 721, 725,		
	to International Patent Classification (IPC) or to be	oth national classification and IPC	
B. FIE	LDS SEARCHED		
Minimum	documentation searched (classification system follow	wed by classification symbols)	
U.S. :	514/284, 356, 460, 568, 574, 721, 725,		
Documenta	tion searched other than minimum documentation to t	he extent that such documents are included	in the fields searched
Electronic	data base consulted during the international search	name of data base and, where practicable	e. search terms used)
1	GISTRY, HCAPLUS, WPIDS, BIOSIS, MEDLIN		<u>-</u>
C. DOC	UMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where	appropriate, of the relevant passages	Relevant to claim No.
Y	Database HCAPLUS on STN (C 118:240454, WO 9302657 A1 (DUR	columbus, OH, USA), No. ANTON, A.), abstract.	1-8, 18-26
Y	Database HCAPLUS on STN (C 127:322650, JP 09268115 A2 (HAM	olumbus, OH, USA), No. ADA, K.), abstract.	1-8, 18-26
Y	CAPLUS on STN (Columbus, OH, 2689008 A1 (BENELLI, L.), abstrac	USA), No. 120:279845, FR t.	1-26
Y	Database HCAPLUS on STN, (C 125:308673, WO 9629045 A1 (DASC	Columbus, OH, USA), No. CALU, A.), abstract.	9-17
X Furth	er documents are listed in the continuation of Box	C. See patent family annex.	
* Spe	ecial categories of cited documents:	"T" later document published after the inter	national filing date or priority
"A" doc	nument defining the general state of the art which is not considered of particular relevance	date and not in conflict with the applic the principle or theory underlying the	invention
	lier document published on or after the international filing date	"X" document of particular relevance; the considered novel or cannot be considered	claimed invention cannot be
cite	nument which may throw doubts on priority claim(s) or which is d to establish the publication date of another citation or other	when the document is taken alone	
"O" doc	cial reason (as specified) ument referring to an oral disclosure, use, exhibition or other	"Y" document of particular relevance; the considered to involve an inventive combined with one or more other such	step when the document is documents, such combination
"P" doe	ans  ument published prior to the international filing date but later than priority date claimed	being obvious to a person skilled in the  "&"  document member of the same patent i	
	actual completion of the international search	Date of mailing of the international sear	rch report
04 OCTO		0.4.11011	1999
Commission Box PCT	nailing address of the ISA/US er of Patents and Trademarks D.C. 20231	Authorized officer	slin for
Facsimile No		Telephone No. (703) 308-1235	V

# INTERNATIONAL SEARCH REPORT

International application No. PCT/US99/16866

		7037771000		
C (Continua	tion). DOCUMENTS CONSIDERED TO BE RELEVANT			
Category*	Citation of document, with indication, where appropriate, of the relevant passages		Relevant to claim N	
Y	Database HCAPLUS on STN, (Columbus, OH, USA), No. 121:308342, JP 06234618 A2 (HIROTA, O.), abstract.	9-17		
Y	Database HCAPLUS on STN, (Columbus, OH, USA), No. 124:185159, WO 6901632 A1 (BROWN, S.), abstract.	1-8, 18-26		
Y	FR 2 666 015 A1 (HOURSEAU, C.) 28 February 1992, pag	9-17		
Y	US 5,650,145 (D. SAINT-LEGER) 22 July 1997, column 2.	9-17		
X	US 4,192,863 T. KONDO) 11 March, 1980, column 4.		27-33.	
			·	
		·	,	
		1		